



INTERNATIONAL ANIMAL SEMEN BANK, INC.
dba INTERNATIONAL CANINE SEMEN BANK and ICSB
MAIN OFFICE | P.O. Box 123 | 430 W Arlington St | Gladstone, Oregon 97027
Phone: 503-663-7031 | FAX: (503) 676-8025
Email: contact@icsb.com | Website: www.ICSB.com

ICSB SU Buffer

When Urine is Present in Canine Semen

In cases where urine is seen in the canine ejaculate, the ICSB SU Buffer can be used. ICSB recommends this buffer for use in all ejaculates that contain urine, whether it will be fresh-collected or fresh-chilled.

Keep the buffer frozen until just prior to use. When needed, thaw the buffer at room to body temperature (70-100°F). It is best to thoroughly and gently mix the buffer with the ejaculate as soon as possible for the best results. If the stud regularly has urine in his sample, we recommend placing the buffer in the collection tube prior to collection for immediate buffering.

Recommend use within 6 months of ship date.

Please contact ICSB if you have any questions or concerns!



INTERNATIONAL ANIMAL SEMEN BANK, INC.
dba INTERNATIONAL CANINE SEMEN BANK and ICSB
MAIN OFFICE | P.O. Box 123 | 430 W Arlington St | Gladstone, Oregon 97027
Phone: 503-663-7031 | FAX: (503) 676-8025
Email: contact@icsb.com | Website: www.ICSB.com

ICSB SU Buffer

When Urine is Present in Canine Semen

In cases where urine is seen in the canine ejaculate, the ICSB SU Buffer can be used. ICSB recommends this buffer for use in all ejaculates that contain urine, whether it will be fresh-collected or fresh-chilled.

Keep the buffer frozen until just prior to use. When needed, thaw the buffer at room to body temperature (70-100°F). It is best to thoroughly and gently mix the buffer with the ejaculate as soon as possible for the best results. If the stud regularly has urine in his sample, we recommend placing the buffer in the collection tube prior to collection for immediate buffering.

Recommend use within 6 months of ship date.

Please contact ICSB if you have any questions or concerns!